

Product Name: Sundew BattleaxePRO Roach Bait
APVMA Approval No: 82708/138943



Label Name:	Sundew BattleaxePRO Roach Bait
Signal Headings:	READ SAFETY DIRECTIONS
Constituent Statements:	0.5 g/kg FIPRONIL
Mode of Action:	
Statement of Claims:	A crevice, crack or spot treatment for the control of cockroaches.
Net Contents:	5 g-100 g
Restraints:	Do not place in areas that are routinely washed.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	
Trade Advice:	
General Instructions:	GENERAL INSTRUCTIONS To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and depress plunger slightly. Recap the dispenser after treatment is completed and return cartridge to outer pack. Re-apply according to remaining level of infestation, when

bait is no longer visibly present. A visual inspection of gel placements is recommended one month after initial treatment.

Resistance Warning:

Precautions:

PRECAUTION
DO NOT apply directly to food, food utensils, food packaging or food preparation surfaces.

Protections:

PROTECTION OF LIVESTOCK, WILDLIFE AND OTHERS
DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. DO NOT apply this product in any areas easily accessible to children or animals.

Storage and Disposal:

STORAGE AND DISPOSAL
Store in the closed original container in a cool, dry place, out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

Safety Directions:

SAFETY DIRECTIONS
May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

First Aid Instructions:

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

First Aid Warnings:

DIRECTIONS FOR USE

Do not place in areas that are routinely washed.

Situation	Pest	Rate	Critical comments
Domestic, commercial and public service buildings	German and other common cockroaches	Apply 1-3 spots per m ² . Recommended spot size is 0.03-0.06 g. Increase the number of spots (up to 3) and spot size depending upon severity of infestation and species present.	Gel placement should be at or near areas where cockroaches gather, such as in cracks and crevices.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION